

New Stability Test For 2-D Systems

Nikos E. Mastorakis

Hellenic Naval Academy, Chair of Computer Science, Terma Hatzikyriakou, 18539, Piraeus, GREECE
Email: mastor@softlab.ntua.gr

Abstract

In this paper, a new stability test for two-dimensional (2-D) discrete systems is presented. The proposed test is based on a simple 1-D test which is simpler but equivalent to that of Schur-Cohn. The test can be easily automated via a suitable computer code. Examples are given.

1. Introduction

2-D systems theory has found applications in a number of areas, for example image enhancement, digital image processing, computerised tomography, remote control, modern circuit design and so on, [4], [6]. Many mathematical problems in 2-D systems theory have already attracted increasing attention in the areas of analysis, synthesis, stability, factorization, controllability, observability, minimality, feedback control and filter design [6].

Stability of two-dimensional (2-D) systems almost arises in all the above applications. Excellent overviews of the stability problem and of the theorems and the tests associated with it can be found in [4÷6]. Some of these theorems check if some 1-D or 2-D polynomials is devoid of zeros in appropriate regions of C (the set of the complex numbers) or $C \times C$ respectively.

A shift-invariant causal single-input single output (SISO) 2-D system can be described by the transfer function:

$$G(z_1, z_2) = \frac{A(z_1, z_2)}{B(z_1, z_2)} \quad (1)$$

where $A(z_1, z_2)$ and $B(z_1, z_2)$ are coprime polynomials in the independent complex variables z_1 and z_2 . It is assumed that there are no nonessential singularities of the second kind on the closed unit bidisk [4], [5], i.e. there are no points (z_1, z_2) with

$$|z_1| \leq 1 \quad \text{and} \quad |z_2| \leq 1 \quad \text{such that} \quad A(z_1, z_2) = B(z_1, z_2) = 0.$$

It is well known that the system (1) is (Hurwitz) stable, [4÷6], if and only if

$$B(0, z_2) \neq 0, \quad \text{for} \quad |z_2| \leq 1 \quad (2.1)$$

$$B(z_1, z_2) \neq 0, \quad \text{for} \quad |z_1| \leq 1, \quad |z_2| = 1 \quad (2.2)$$

Additionally, the polynomial $B(z_1, z_2)$ is said to be *VSHP* (Very Strict Hurwitz Polynomial) if and only if (2.1) and (2.2) are fulfilled. Condition (2.1) is relatively easy to check using any 1-D stability test. Condition (2.2) is more difficult since it includes two variables. For Condition (2.2), several tests already exist which check various conditions of the coefficients of $f(z_1, z_2)$, [1], [3], [4÷7], [11]. We denote

$$B(z_1, z_2) = \sum_{i=0}^{n_1} b_i(z_2) z_1^i \quad (3)$$

By interchanging the variables z_1 and z_2 , other Conditions symmetrical to (2.1) and (2.2) are found.

In this paper, a new test for checking Condition (2.2) is proposed. The analysis is based on a result concerning 1-D systems stability [2]. The theoretical results are illustrated by two examples.

2 Main Results

Let the degree in z_1 and z_2 of $B(z_1, z_2)$ be n_1 and n_2 respectively. Then $B(z_1, z_2) \neq 0$, for $|z_1| \leq 1$, $|z_2| = 1$ if and only if

$B_1(z_1, z_2) \neq 0$, for $|z_1| \geq 1$, $|z_2| = 1$ where

$B_1(z_1, z_2) = z_1^{n_1} B(z_1^{-1}, z_2)$. The polynomial

$B_1(z_1, z_2)$ is written as a polynomial in z_1 with coefficients which are polynomials in z_2 :

$$B_1(z_1, z_2) = \sum_{i=0}^{n_1} b_{1,i}(z_2) z_1^i \quad (4)$$

with $b_{1,i}(z_2) = b_{n_1-i}(z_2)$ and $i = 0, \dots, n_1 - 1$

The following Theorem summarises a result obtained by Barnett [2]

Theorem 1: The polynomial

$a(z) = z^n + a_{n-1}z^{n-1} + \dots + a_1z + a_0$ does not have

any zeros outside the unit circle $|z| = 1$ if and only if

the matrix $C = \bar{a}_0 A^n + \bar{a}_1 A^{n-1} + \dots + \bar{a}_{n-1} A + I_n$ (the bar indicates complex conjugate) is positive definite, where

$$A = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ & & \vdots & & \\ 0 & 0 & 0 & \dots & 1 \\ -a_0 & -a_1 & -a_2 & \dots & -a_{n-1} \end{bmatrix} \quad (5)$$

and I_n is the $n \times n$ identity matrix.

Based on this Theorem, we can formulate a Test for checking Condition (2.2). The Test is summarised by the following Theorem:

Theorem 2: $B(z_1, z_2) \neq 0$, for $|z_1| \leq 1$, $|z_2| = 1$, if and only if the matrix

$$C(z_2) = \bar{a}_0(z_2) A^{n_1} + \bar{a}_1(z_2) A^{n_1-1} + \dots + \bar{a}_{n_1-1}(z_2) A + I_{n_1}$$

is positive definite for all z_2 with $|z_2| = 1$, where

$$A = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ & & \vdots & & \\ 0 & 0 & 0 & \dots & 1 \\ -a_0(z_2) & -a_1(z_2) & -a_2(z_2) & \dots & -a_{n_1-1}(z_2) \end{bmatrix} \quad (6)$$

and $a_i(z_2) = \frac{b_{1,i}(z_2)}{b_{1,n_1}(z_2)}$ or equivalently

$$a_i(z_2) = \frac{b_{n_1-i}(z_2)}{b_0(z_2)} \quad (7)$$

$i = 0, \dots, n_1 - 1$.

Proof: It follows directly from Theorem 1.

Furthermore, we state another theorem which is an improvement of the Theorem 2.

Theorem 3: $B(z_1, z_2) \neq 0$, for $|z_1| \leq 1$, $|z_2| = 1$, if and only if the matrix

$$C(z_2) = \bar{a}_0(z_2) A^{n_1} + \bar{a}_1(z_2) A^{n_1-1} + \dots + \bar{a}_{n_1-1}(z_2) A + I_{n_1}$$

(i) is positive definite for a z_0 , with $|z_0| = 1$,

and (ii) $\det\{C(z_2)\} > 0$ for all z_2 with $|z_2| = 1$,

where A is provided by (6) and (7).

Proof: It was pointed out by Siljak, [12], that for positivity checking of such a matrix, one requires the positivity checking of the matrix at one point, say at

z_0 , with $|z_0| = 1$ and the positivity checking of the

determinant for all z_1 with $|z_1| = 1$.

Remark 1: The above Theorem provides a Test which constitutes a simplification, since it does not require the matrix $C(z_2)$ to be positive definite for all

z_2 , with $|z_2| = 1$.

Remark 2: One should notice that the matrix C is easily evaluated using the Horner formula as follows

$$C = I_n + \dots + A(\bar{a}_2 I + A(\bar{a}_1 I + \bar{a}_0 A)) \quad (8)$$

Taking into account the special structure of A , for the computation of C , we find a total cost of $n + (3n - 2)(n - 1)$ multiplications and $n + (3n - 2)(n - 1)$ additions i.e. $O(3n^2)$ multiplications and $O(3n^2)$ additions. The method of Schur-Cohn includes $n^2(n + 1)$ multiplications i.e. $O(n^3)$ multiplications and $n^2(n + 1)/2$ additions i.e. $O(n^3/2)$ additions. Therefore, the present method is better than the classical method of Schur-Cohn with respect to the computational complexity.

Example 1: We consider the Example given in [4]

$$B(z_1, z_2) = (12 + 6z_2) + (10 + 5z_2)z_1 + (2 + z_2)z_1^2 \quad (9)$$

Obviously, $B(0, z_2) \neq 0$, for $|z_2| \leq 1$. Therefore Condition (2.1) holds. In order to check Condition (2.2), we form the polynomial:

$$B_1(z_1, z_2) = (2 + z_2) + (10 + 5z_2)z_1 + (12 + 6z_2)z_1^2 \quad (10)$$

So, we obtain

$$A = \begin{bmatrix} 0 & 1 \\ -\frac{2 + z_2}{12 + 6z_2} & -\frac{10 + 5z_2}{12 + 6z_2} \end{bmatrix} \quad (11)$$

Now, one has

$$C(z_2) = I_2 + A \left(-\frac{10 + 5\bar{z}_2}{12 + 6\bar{z}_2} - \frac{2 + \bar{z}_2}{12 + 6\bar{z}_2} A \right) \quad (12)$$

After the necessary algebraic manipulation, one finds

$$C(z_2) = \begin{bmatrix} \frac{35}{36} & \frac{25}{36} \\ -\frac{25}{216} & \frac{85}{216} \end{bmatrix} \quad (13)$$

Clearly, this matrix is positive definite. Therefore the 2-D (discrete) system having the characteristic polynomial $B(z_1, z_2)$ is stable. The result agrees with that in [4].

Example 2: Consider the Example given in [9].

$$B(z_1, z_2) = (13 - z_2) + (-12 - 7z_2)z_1 + 8z_1^2 \quad (14)$$

Since $B(0, z_2) \neq 0$, for $|z_2| \leq 1$, Condition (2.1) clearly holds. In order to check Condition (2.2), we consider the polynomial:

$$B_1(z_1, z_2) = 8 + (-12 - 7z_2)z_1 + (13 - z_2)z_1^2 \quad (15)$$

So, we obtain

$$A = \begin{bmatrix} 0 & 1 \\ -\frac{8}{13 - z_2} & \frac{12 + 7z_2}{13 - z_2} \end{bmatrix} \quad (16)$$

Then

$$C(z_2) = I_2 + A \left(-\frac{12 + 7\bar{z}_2}{13 - \bar{z}_2} + \frac{8}{13 - \bar{z}_2} A \right) \quad (17)$$

For convenience, if one sets $z_2 = e^{j0.5}$, one has only numerical manipulation and finally

$$C(e^{j0.5}) = \begin{bmatrix} 0.5652 & -0.4972 + j0.5179 \\ 0.3411 - j0.3283 & -0.3462 + j0.6014 \end{bmatrix} \quad (18)$$

The determinant of this matrix is $-0.1960 < 0$. Evidently, this matrix is negative definite and the corresponding 2-D (discrete) system is unstable. The result agrees with that of [9].

3 Conclusion

The test and its implementation is based on a result published by Barnett and is easy to use. Note that if one interchanges the role of the variables z_1 and z_2 , another equivalent form of the method can be presented. Work is in progress by the author in the area of 2-D stability theorems formulating new methods for 2-D stability margin [10] and necessary conditions for 2-D systems stability [9].

References

1. Anderson B.D.O. and Jury E.I., "Stability test for two-dimensional recursive filters", *IEEE Tr. Aud Electroacoust.*, **20**, (1972) 366-372,
2. Barnett S., "Number of zeros of a complex polynomial inside the unit circle", *Electronics Letters*, **6**, 6, (1970), 164-165.
3. Huang T.S., "Stability of two-dimensional recursive filters", *IEEE Trans. Audio Electroacoust.*, **20**, (1972), 158-163, .
4. Jury, E.I., "Stability of multidimensional systems and related problems", *Multidimensional Systems - Techniques and Applications*, S.G. Tzafestas editor, N. York :Marcell Dekker, 1985
5. Jury, E.I., "Stability of multidimensional scalar and matrix polynomials", *Proceedings of the IEEE*, **66**, 9, (1978), 1018-10476.
6. Kaczorek T., *2-D Linear Systems*, Springer-Verlang, Lecture Notes in Control and Information Sciences, Berlin-Heidelberg, 1985.
7. Maria G.A. and Fahmy M.M., "On the stability of two-dimensional digital filters", *IEEE Tr. Audio Electroacoust.*, **21**, (1973), 470-472.
8. Mastorakis N.E., "Comments Concerning Multidimensional Polynomials Properties" *IEEE Tr..Autom. Cntrol*, **41**, No.2, (1996).
9. Mastorakis N. E., "New Necessary Stability Conditions for 2-D Systems", *IEEE Trans. Circuits & Systems-I*: (1994-...under review).
10. Mastorakis N.E. "Stability Margin of Two-Dimensional Continuous Systems" *IEEE Trans.Autom. Control*, (submitted).
11. Shanks J.L., Treital S., and Justice J.H., "Stability and synthesis of two dimensional recursive filters", *IEEE Trans. Audio Electroacoust.*, **20**, (1972), 115-208.
12. Siljak, "Stability criteria for two-variable polynomials", *IEEE Trans. Circuits Syst.*, **22**, (1975) 183-189.

Nikos E. Mastorakis: received his B.Sc.-M.Sc. (Diploma) in Electrical Engineering from the National Technical University of Athens in 1988 and the Ph.D. in Electrical Engineering and Computer Science from the same university in 1992. In 1995 he also received the B.Sc. (Ptychion) in Pure Mathematics from the National University of Athens. From 1993 to 1994 he served his obligatory military service in the Greek Army as a special scientist on Computers and Electronics. From 1988 to 1995 he taught courses in the Electrical Engineering Department of the National Technical University of Athens. He is currently Professor in the Hellenic Naval Academy (which is considered under the Greek law as a Military University ("Anotato Ekpaideutiko Idryma" of the Hellenic Ministry of Defence) equivalent to the other greek civilian universities).

Dr. Mastorakis directs the Chair of Computer Science in the H.N.Academy. He is an active researcher in the areas of *applied mathematics (systems theory) and computer science*. He is the author of 5 books in the Greek literature and the author of more than 70 papers in international journals and conferences. He has received several awards for his academic studies and his scientific research. He is the Organizer and the Chairman of the international conference on *Circuits, Systems and Computers*. Dr. Mastorakis is a member of New York Academy of Sciences, of A.F. Communications and Electronics Association and other scientific societies and a registered professional electrical and mechanical engineer. His biography is included in international biographical reference publications (Marquis Who is Who in the World, 1998)