

F. COUENNE-CELLE, H. HAMMOURI, J.-C. MARQUES

mailing address : L.A.G.E.P, URA CNRS D1328 bâtiment 721 troisième étage
43 Boulevard du 11 Novembre 1918
69622 VILLEURBANNE CEDEX
FRANCE

e-mail : couenne@chreon.univ-lyon1.fr
phone number : 72 43 11 94 or 72 43 11 85

Keywords : nonlinear dissipative systems, observers, separation principle, feedback stabilization.

In this paper, we consider a multi input nonlinear system of the form

$$(\Sigma) \quad \begin{cases} \dot{x} = f(x) + \sum_{i=1}^m u_i(t) g_i(x) \\ y = h(x) \end{cases}$$

with $x(t) \in \mathbb{R}^n$, $y(t) \in \mathbb{R}$, $u(t) \in \mathbb{R}^m$. (Σ) is supposed to be dissipative, that is to say there exists a C^∞ function $V : \mathbb{R}^n \rightarrow \mathbb{R}^+$ verifying the following properties :

$$(P.1) \quad \lim_{\|x\| \rightarrow +\infty} V(x) = +\infty ; V(0) = 0.$$

$$(P.2) \quad L_f(V)(x) \leq 0, \forall x \in \mathbb{R}^n$$

In the case where $L_f(V) = 0$ the system is said to be conservative.

The aim is to show that if this class of systems are stabilizable by some feedback control $u(x)$ then it can be performed by some feedback $u(\hat{x})$ where \hat{x} is an estimate of x coming from an observer.

For some particular classes of nonlinear systems, this problem has been treated in [TS], [HA, BU].

Here we present an extension of the bilinear dissipative separation principle proposed in [GA, KU].

First we recall the stabilization result for (Σ) .

This result has been already proved for conservative nonlinear systems in [GA, BO], and for bilinear dissipative systems in [GA, KU]. The result is the following :

Theorem 1 :

Assume that (Σ) satisfies :

a : there exists a real valued function V with $V(0) = 0$ satisfying (P.1) and (P.2)

b : $dV(x) = 0 \iff x = 0$

c : $f(0) = 0$

d : the vector space E_x spanned by $\{f(x), ad_f^k(g_i)(x), k \in \mathbb{N}, i = 1, \dots, m\}$ is n -dimensional, for any x such that $L_f(V)(x) = 0, L_{g_i}(V)(x) = 0$ and $x \neq 0$.

Then the feedback $u(x) = \begin{pmatrix} u_1(x) \\ \vdots \\ u_m(x) \end{pmatrix}$ where $u_i(x) = -r_i(x)L_{g_i}(V)(x)$ stabilizes (Σ) , and that for any C^∞ function $r_i(x), r_i(x) > 0 \forall x \in \mathbb{R}^n \setminus \{0\}$.

The proof of this theorem uses the same steps as in [GA,BO].

Before stating the main theorem, we need the following assumptions :

(P.3) There exists an observer \mathcal{O} for (Σ) :

$$\begin{cases} \dot{\hat{x}} = f(\hat{x}) + \sum_{i=1}^m u_i g_i(\hat{x}) + K(t)(h(\hat{x}) - y) \\ \dot{K}(t) = P(K(t), u(t), y(t), \hat{x}(t)) \quad \text{with } \hat{x} \in \mathbb{R}^n, \end{cases}$$

\mathcal{O} converges exponentially for any input $u \in L^\infty(\mathbb{R}), \|u\|_{L^\infty} \leq r$ and the gain $K(t)$ is uniformly bounded (Note that the boundedness of the gain $K(t)$ is always satisfied in the litterature).

(P.4) The system (Σ) satisfies the conditions of theorem 1 and the $r_i(x)$'s are chosen such that $|u_i(x)| \leq r$.

Consider the following system :

$$(G) \quad \begin{cases} \dot{x} = f(x) + \sum_{i=1}^m u_i(x)g_i(x) \quad \text{with } u_i(x) = -r_i(x)L_{g_i}V(x) & (1) \\ \dot{\hat{x}} = f(\hat{x}) + \sum_{i=1}^m u_i g_i(\hat{x}) + K(t)(h(\hat{x}) - h(x)) & (2) \\ \dot{K}(t) = P(K(t), u(t), y(t), \hat{x}(t)) & (3) \end{cases}$$

We obtain the main result :

Theorem 2 (separation principle) :

With the hypotheses (P.3) and (P.4), if, moreover :

(P.5) For any compact K of \mathbb{R}^n , there exists a compact K' ($K \subset K'$) such that for any trajectory of (G) which are issued from $K \times K \times \mathbb{R}^n$, its components $x(t)$ and $\hat{x}(t)$ lie in $K' \times K'$.

Then the set $\{0\} \times \{0\} \times \mathbb{R}^n$ is an attractor of any trajectory of (G) issued from $K \times K \times \mathbb{R}^n$. Moreover the subsystem formed by (1) and (2) is globally asymptotically stable.

In what follows, we give a sufficient condition for which the system (G) satisfies (P.5). For this, recall that a Lyapunov function for an autonomous system $\dot{x} = F(x)$ at the equilibrium point 0 is a C^1 function :

$W : \mathbb{R}^n \longrightarrow \mathbb{R}^+$ such that :

i) $W(x) = 0 \iff x = 0$

ii) $\lim_{x \rightarrow \infty} W(x) = +\infty$

iii) $\frac{\delta W(x)}{\delta x}(x) \cdot F(x) < 0, \forall x \neq 0$.

($\frac{\delta W(x)}{\delta x}(x)$ is a one-line vector and $F(x)$ is a one-column vector.)

Remarks :

1) The above function $V(x)$ is not, in general, a Lyapunov function.

2) By J.L. Massera's theorem [MA], any C^1 globally asymptotically stable system admits a Lyapunov function.

Using the same notations as above, and denoting by (Σ_C) the closed loop system $\dot{x} = f(x) + \sum_{i=1}^m u_i(x)g_i(x)$ with $u_i(x) = -r_i(x)L_{g_i}(V(x))$ we have :

Proposition 3 :

The condition (P.5) of theorem 2 holds if one of the following assumption is satisfied :

(A₁) : The above function $V(x)$ verifies : $\|\frac{\partial V}{\partial x}\| \leq \lambda_1(1 + |V(x)|)$, $\forall x \in \mathbb{R}^n$, $\lambda_1 > 0$

(A₂) :
 i) There exists a Lyapunov function $W(x)$ for Σ_C which satisfies :
 $\|\frac{\partial W}{\partial x}\| \leq \lambda_2(1 + |W(x)|)$, $\forall x \in \mathbb{R}^n$, $\lambda_2 > 0$
 ii) The output function $h(x)$ is globally Lipschitzian.

References :

- [HA, DE] H. Hammouri, J. De Leon Morales. "Observer synthesis for state affine systems", Proceeding of the 29th CDC Honolulu, Hawaii, Dec 5-7 1991.
- [BO, CO, CE] G. Bornard, N. Couenne, F. Celle. "Regularly persistent observers for bilinear systems" Proceedings of the 29th international conference on nonlinear systems. New trends in nonlinear systems theory. Lecture notes in control and information sciences, Vol 122, Springer Verlag, June 1988.
- [GA, KU] J.P. Gauthier, I. Kupka. "A separation principle for bilinear systems with dissipative drifts", to appear in IEEE Transactions on Automatic Control.
- [GA, BO] J.P. Gauthier, G. Bornard. "Stabilisation des systèmes non-linéaires", Outils et modèles mathématiques pour l'automatique, l'analyse des systèmes et le traitement du signal, Edition du CNRS, 1983.
- [MA] J.L. Massera. "Contribution to stability theory", Ann. of Math., 64 (1956), 182-206.
- [HA, BU] H. Hammouri, G. Busawon. "A global stabilization of a class of nonlinear systems by means of an observer", To appear in Applied Mathematics Letters
- [TS] Tsiniias. "A theorem on global stabilization of nonlinear systems by linear feedback", Systems and Control Letters, 17, (1991).