

A Polynomial Approach to the Solution of the Invariant Polynomial Assignment Problem for Periodic Systems *

L. Jetto and S. Longhi

Dipartimento di Elettronica ed Automatica
Università di Ancona, via Brece Bianche, 60131 Ancona, Italy.

Fax: +39 71 2804334,

E-mail: JETTO@ANVAX2.UNIAN.IT

E-mail: LONGHI@ANVAX1.UNIAN.IT

Abstract

This paper considers the problem of assigning the closed loop invariant polynomials of a unity feedback control system, where the plant is a linear, discrete-time, periodic system. By a matrix algebraic approach, the solution is provided in terms of a linear, discrete-time periodic controller whose associate transfer matrix originates from the solution of a suitable diophantine matrix equation. The particular form chosen for this solution guarantees the causality of the periodic realization of the associate transfer matrix.

1 Introduction

Various classes of processes, such as periodically time-varying networks and filters (for example switched-capacitors circuits and multirate digital filters), chemical processes, multirate sampled-data systems, can be modeled through a linear periodic system (see, e.g., [1], [2] and references therein). Moreover, the study of linear periodic systems can be helpful even for the stabilization and control of time-invariant linear systems through a periodic controller, that has been recently investigated [3], [4], [5], [6], [7], [8], and for the stabilization and control of a class of bilinear systems [9], [10], [11].

In the discrete-time case, a control theory is developing with the help of algebraic and geometric techniques and contributions on several control problem have been given, including eigenvalues assignment,

state and output dead-beat control, disturbance decoupling, model matching, adaptive control, robust control and optimal H_2/H_∞ control (see, e.g., [2], [12], [13], [14], [15], [16], [17], [18], [19], [20], [21], [22], [23], [24]).

The aim of this paper is to analyze the invariant polynomial assignment problem for the class of discrete-time linear periodic systems. This problem generalizes the characteristic polynomial assignment, which, for the same class of systems, was solved by a geometric approach in [17], [21], [23]. For time-invariant plants, the invariant polynomial assignment was considered in [4], [7], [25], [26].

The paper is organized in the following way. In Section 2 preliminary definitions and results are given. The problem considered in this paper is formally stated in Subsection 3.1, and conditions for its solvability are constructively established in Subsection 3.2.

2 Preliminary results

Consider the ω -periodic discrete-time system Σ described by

$$x(k+1) = A(k)x(k) + B(k)u(k) \quad (2.1)$$

$$y(k) = C(k)x(k) + D(k)u(k) \quad (2.2)$$

where $k \in \mathbb{Z}$, $x(k) \in \mathbb{R}^n$ is the state, $u(k) \in \mathbb{R}^p$ is the input, $y(k) \in \mathbb{R}^q$ is the output and $A(\cdot), B(\cdot), C(\cdot), D(\cdot)$ are periodic matrices of period ω (briefly, ω -periodic). Denote also by $\Phi(k, k_0)$, $k \geq k_0$, the transition matrix associated with $A(\cdot)$.

It is well-known that, for any initial time $k_0 \in \mathbb{Z}$, the output response of system Σ for $k \geq k_0$, to given

*Work supported by the Ministero dell'Università e della Ricerca Scientifica

initial state $x(k_0)$ and control function $u(\cdot)$, can be obtained through the time-invariant associated system of Σ at time k_0 , denoted by $\Sigma^a(k_0)$ [27]. $\Sigma^a(k)$ is represented by

$$x_k(h+1) = E_k x_k(h) + J_k u_k(h) \quad (2.3)$$

$$y_k(h) = L_k x_k(h) + M_k u_k(h) \quad (2.4)$$

where $E_k := \Phi(\omega + k, k)$, $J_k := [(J_k)_1 \cdots (J_k)_\omega]$, $(J_k)_i := \Phi(\omega + k, i + k)B(i - 1 + k)$, $i = 1, \dots, \omega$, $L_k := [(L_k)_1' \cdots (L_k)_\omega']$, $(L_k)_i := C(i - 1 + k)\Phi(i - 1 + k, k)$, $i = 1, \dots, \omega$, $M_k := [(M_k)_{ij} \in \mathbb{R}^{q \times p}$, $i, j = 1, \dots, \omega]$, with $(M_k)_{ij} := C(i - 1 + k)\Phi(i - 1 + k, j + h)B(j - 1 + k)$, if $i > j$, $(M_k)_{ij} := D(i - 1 + h)$, if $i = j$ and $(M_k)_{ij} := 0$, if $i < j$.

In fact, if $x_k(0) = x(k)$ and $u_k(h) := [u'(h\omega + k) \ u'(h\omega + k + 1) \cdots u'(h\omega + k + \omega - 1)]'$ for all $h \in \mathbb{Z}^+$, then $x_k(h) = x(k + h\omega)$ and $y_k(h) = [y'(h\omega + k) \ y'(h\omega + k + 1) \cdots y'(h\omega + k + \omega - 1)]'$ for all $h \in \mathbb{Z}^+$. The notion of associated system at time k allows one to analyze structural and stability properties and pole-zero-structures of periodic systems [1], [28], [29], [30], [31]. For example, the subspace of reachable (unobservable) states of system Σ at time k is readily seen to coincide with that of system $\Sigma^a(k)$ if it is expressed in terms of matrices E_k , J_k , L_k and M_k [29]. Obviously, $\Sigma^a(k + \omega) = \Sigma^a(k)$ for all integer k .

The notions of invariant zero, transmission zero and pole of the ω -periodic system Σ at time k are defined with reference to the following $\omega q \times \omega p$ matrix

$$W_k(d) = L_k d(I_n - dE_k)^{-1} J_k + M_k, \quad (2.5)$$

where I_n denotes the identity matrix of dimension n and $d := z^{-1}$ is the backward shift operator. The rational matrix $W_k(d)$ is the transfer matrix of the associated system of Σ at time k and is called the *associated transfer matrix of Σ at time k* . A complete analysis of pole-zero structure of system Σ is reported in [29] and [32] making use of the associated transfer matrix characterized with the forward shift operator z . The following result, that follows from Lemma 2.1 in [29], shows the dependence of $W_k(d)$ with respect to the initial time k .

Lemma 2.1 *For any integer k it holds that:*

$$W_{k+1}(d) = \begin{bmatrix} 0 & I_{q(\omega-1)} \\ d^{-1}I_q & 0 \end{bmatrix} W_k(d) \begin{bmatrix} 0 & dI_p \\ I_{p(\omega-1)} & 0 \end{bmatrix}. \quad (2.6)$$

As a consequence of this result the rank r of

$W_k(d)$ is independent of time k (see, e.g., [29] for a similar result with the forward shift operator z).

The transfer matrix $W_k(d)$ can be factored as

$$W_k(d) = A_k^{-1}(d) B_k(d) = \bar{B}_k(d) \bar{A}_k^{-1}(d), \quad (2.7)$$

where $A_k(d)$ and $B_k(d)$ are relatively left prime (*rlp*) polynomial matrices and $\bar{A}_k(d)$ and $\bar{B}_k(d)$ are relatively right prime (*rrp*) polynomial matrices. Moreover, denoting with $T_k(d)$ the Smith-McMillan form of $W_k(d)$ the following relation holds

$$W_k(d) = U_k^{W,L}(d) T_k(d) U_k^{W,R}(d), \quad (2.8)$$

with

$$T_k(d) = \begin{bmatrix} \text{diag}\{\epsilon_1^k(d)/\psi_1^k(d), \dots, \epsilon_r^k(d)/\psi_r^k(d)\} & 0_{r, \omega p - r} \\ 0_{\omega q - r, r} & 0_{\omega q - r, \omega p - r} \end{bmatrix},$$

where $\epsilon_i^k(d)$ and $\psi_i^k(d)$ ($i = 1, \dots, r$) are coprime polynomials such that $\epsilon_i^k(d)$ ($\psi_{i+1}^k(d)$) divides $\epsilon_{i+1}^k(d)$ ($\psi_i^k(d)$), and $U_k^{W,L}(d)$ and $U_k^{W,R}(d)$ are unimodular matrices. The $\epsilon_i^k(d)$'s and $\psi_i^k(d)$'s ($i = 1, \dots, r$) are uniquely determined by $W_k(d)$ up to arbitrary real scalars. Analogously to the time-invariant case [25], the polynomials $\psi_i^k(d)$ ($i = 1, \dots, r$) are called the *invariant polynomials of Σ at time k* . As shown in [29], [32], under the hypothesis of reachability and observability of Σ at all times, the product of these polynomials characterizes the stability properties of Σ .

By (2.8), the *rlp* polynomial matrices $A_k(d)$ and $B_k(d)$ and the *rrp* polynomial matrices $\bar{A}_k(d)$ and $\bar{B}_k(d)$ satisfying (2.7) are given by

$$A_k(d) = [\text{diag}\{\psi_1^k(d), \dots, \psi_r^k(d)\} \ I_{q-r}] (U_k^{W,L})^{-1}, \quad (2.9)$$

$$B_k(d) = [\text{diag}\{\epsilon_1^k(d), \dots, \epsilon_r^k(d)\} \ 0_{q-r, p-r}] U_k^{W,R}, \quad (2.10)$$

$$\bar{A}_k(d) = (U_k^{W,R})^{-1} [\text{diag}\{\psi_1^k(d), \dots, \psi_r^k(d)\} \ I_{q-r}], \quad (2.11)$$

$$\bar{B}_k(d) = U_k^{W,L} [\text{diag}\{\epsilon_1^k(d), \dots, \epsilon_r^k(d)\} \ 0_{q-r, p-r}]. \quad (2.12)$$

The above definition of invariant polynomials of Σ at time k and equations (2.7), (2.8), (2.9), (2.10), (2.11) and (2.12) yield the following result. (Note that, two polynomials are called associate if their ratio is a scalar [25].)

Lemma 2.2 *For any integer k , the invariant polynomials of Σ at time k are associate of the invariant polynomials of the Smith forms of $A_k(d)$ and $\bar{A}_k(d)$.*

Denote by $\chi(p, q, \omega)$ the class of $\omega q \times \omega p$ rational matrices

$$W(d) = \begin{bmatrix} W_{11}(d) & W_{12}(d) & \cdots & W_{1\omega}(d) \\ W_{21}(d) & W_{22}(d) & \cdots & W_{2\omega}(d) \\ \vdots & \vdots & \ddots & \vdots \\ W_{\omega 1}(d) & W_{\omega 2}(d) & \cdots & W_{\omega\omega}(d) \end{bmatrix},$$

$$W_{ij}(d) \in \mathbb{C}^{q \times p}, i, j = 1, \dots, \omega, \quad (2.13)$$

with $W_{ij}(0) = 0, i < j, i, j = 1, \dots, \omega$. The class $\chi(q, p, \omega)$ characterizes the transfer matrices of ω -periodic systems. In fact, the causality of ω -periodic system Σ implies that the associated transfer matrix of Σ at time k belongs to the class $\chi(p, q, \omega)$ for all $k \in \mathbb{Z}$ [33]. Then, the causality of Σ implies that the roots of invariant polynomials of Σ at time k are different to zero for all integer k . This in turn implies that matrices $A_k(0)$ and $\bar{A}_k(0)$ are nonsingular. Foregoing considerations and Lemma 2.1 allow us to prove the following result.

Lemma 2.3 *The invariant polynomials of Σ at time k are independent of k .*

Moreover, $\chi(p, q, \omega)$ characterizes also the class of rational matrices that can be realized by an ω -periodic system of the form (2.1), (2.2). The solution of the minimal realization problem for the periodic case is described by a system reachable and observable at any time whose matrices have generally time-varying dimensions. In general, the subspaces of reachable states and/or observable states may have time-varying dimensions. Therefore, it is natural, in order to consistently solve the problem of minimal realization, to allow for state-space description having time-varying dimensions. The possibility of computing a "quasi" minimal (reachable and observable at least in one time) uniform (fixed-dimension) realization is also available. Efficient algorithms for the computation of minimal or quasi minimal realization of a given transfer matrix are introduced in [33] and [34].

Moreover, the associated system at a given time k of a composite system obtained connecting ω -periodic subsystems coincides with the same connection of the associated systems at time k of the composing subsystems [32].

3 Main result

3.1 Problem statement

Consider an ω -periodic controller Σ_G for system Σ acting in an unity feedback error actuated servo sys-

tem and described by

$$x_G(k+1) = A_G(k)x_G(k) + B_G(k)e(k), \quad (3.1)$$

$$u(k) = C_G(k)x_G(k) + D_G(k)e(k), \quad (3.2)$$

where $e(k) := r(k) - y(k)$, $r(\cdot)$ being the external reference to be tracked, and $x_G(k) \in \mathbb{R}^{n_G}$ is the state.

The problem considered in this paper is formally stated as follows.

Problem 3.1 Given an ω -periodic system Σ reachable and observable at all times, and m causal polynomials $s_1(d), s_2(d), \dots, s_m(d)$ such that $s_{i+1}(d)$ divides $s_i(d)$, find an ω -periodic controller Σ_G described by (3.1), (3.2), such that the closed loop system be minimally realized and its invariant polynomials be associated of $s_i(d), i = 1, 2, \dots, m$.

Note that, by Lemma 2.3, Problem 3.1 has been stated independently of the time instant k .

The $\omega p \times \omega q$ associated transfer matrix of Σ_G at time k is expressed by

$$G_k(d) = L_k^G d(I_{n_G} - dE_k^G)^{-1} J_k^G + M_k^G, \quad (3.3)$$

where matrices $L_k^G \in \mathbb{R}^{\omega p \times n_G}, E_k^G \in \mathbb{R}^{n_G \times n_G}, J_k^G \in \mathbb{R}^{n_G \times \omega q}$ and $M_k^G \in \mathbb{R}^{\omega p \times \omega q}$ are defined as matrices L_k, E_k, J_k and M_k with matrices $A(\cdot), B(\cdot), C(\cdot)$ and $D(\cdot)$ substituted respectively by matrices $A_G(\cdot), B_G(\cdot), C_G(\cdot)$ and $D_G(\cdot)$.

The causality of system Σ_G implies that $G_k(d)$ belongs to the class $\chi(p, q, \omega)$.

Let $G_k(d)$ be factored as

$$G_k(d) = P_k^{-1}(d) Q_k(d) \quad (3.4)$$

where $P_k(d)$ and $Q_k(d)$ are rlp polynomial matrices.

3.2 Problem solution

The solvability condition of the problem considered is stated in the following theorem.

Theorem 3.1 *Problem 3.1 has a solution if and only if $m \leq \min(\omega q, \omega p)$.*

Proof (Necessity) Let $W_k^C(d)$ be the $\omega q \times \omega q$ associated transfer matrix at time k of the ω -periodic closed loop system Σ_C described by (2.1), (2.2), (3.1), (3.2), and let it be factored as

$$W_k^C(d) = C_k^{-1}(d) D_k(d) = \bar{D}_k(d) \bar{C}_k^{-1}(d). \quad (3.5)$$

where $C_k(d)$ and $D_k(d)$ are rlp polynomial matrices and $\bar{D}_k(d)$ and $\bar{C}_k(d)$ are rrp polynomial matrices.

By Lemma 2.2 applied to the closed loop system, the invariant polynomials of Σ_C at time k are associated to the invariant polynomials of the Smith forms of $C_k(d)$ and $\bar{C}_k(d)$. Moreover, taking into account that the closed loop system Σ_C is required to be free of hidden modes, it follows that the invariant polynomials of Σ_C are also associated to the invariant polynomials of the $\omega p \times \omega p$ polynomial matrix $X_k(d)$ such that [25]

$$P_k(d)\bar{A}_k(d) + Q_k(d)\bar{B}_k(d) = X_k(d). \quad (3.6)$$

As the invariant polynomials of the $\omega p \times \omega p$ polynomial matrix $X_k(d)$ are associated to the invariant polynomials of the $\omega q \times \omega q$ polynomial matrices $C_k(d)$ and $\bar{C}_k(d)$, it follows that the number m of invariant polynomials that can be assigned can not be larger than $\min(\omega p, \omega q)$.

(Sufficiency) By the primeness of the pair $(\bar{B}_k(d), \bar{A}_k(d))$, equation (3.6), can be solved with respect to $P_k(d)$, $Q_k(d)$, for any choice of $X_k(d)$. The matrix $Q_k(d)$ will be sought of the form $Q_k(d) = dQ_k^a$ because, as it will be shown later, this guarantees the causality of the ω -periodic controller Σ_G . With this choice of $Q_k(d)$, equation (3.6), can be rewritten as

$$P_k(d)\bar{A}_k(d) + dQ_k^a(d)\bar{B}_k(d) = X_k(d). \quad (3.7)$$

By the causality of Σ , one has that $\det \bar{A}_k(0) \neq 0$, so that right primeness of $\bar{A}_k(d)$ and $\bar{B}_k(d)$ implies right primeness of $\bar{A}_k(d)$ and $d\bar{B}_k(d)$; this in turn implies that also equation (3.7) can be solved with respect to $P_k(d)$ and $Q_k^a(d)$ for each $X_k(d)$. The general solution of (3.7) is of the form

$$\begin{bmatrix} P_k(d) & Q_k^a(d) \end{bmatrix} = \begin{bmatrix} X_k(d) & T(d) \end{bmatrix} \begin{bmatrix} L_2(d) & R_2(d) \\ M_2(d) & S_2(d) \end{bmatrix}, \quad (3.8)$$

where $T(d)$ is an arbitrary polynomial matrix and where the matrix

$$U(d) := \begin{bmatrix} L_2(d) & R_2(d) \\ M_2(d) & S_2(d) \end{bmatrix}$$

is a unimodular matrix whose elements are polynomial matrices such that

$$\begin{aligned} L_2(d)\bar{A}_k(d) + M_2(d)d\bar{B}_k(d) &= I_p, \\ R_2(d)\bar{A}_k(d) + S_2(d)d\bar{B}_k(d) &= 0. \end{aligned}$$

Unimodularity of $U(d)$ implies that the matrix $\begin{bmatrix} P_k(d) & Q_k^a(d) \end{bmatrix}$ is full row rank for all $d \in \mathbb{C}$, if and only if the same property holds for $\begin{bmatrix} X_k(d) & T(d) \end{bmatrix}$. As $T(d)$ is arbitrary, this last requirement can always be satisfied, so that $P_k(d)$ and $Q_k^a(d)$ solution

of (3.7) are *rlp* polynomial matrices. Nonsingularity of $X_k(0)$ and $\bar{A}_k(0)$ implies nonsingularity of $P_k(0)$, so that also $P_k(d)$ and $Q_k(d) = dQ_k^a(d)$, solutions of (3.6), are *rlp* polynomial matrices.

The transfer matrix $G_k(k) = P_k(d)^{-1}Q_k(d) = P_k(d)^{-1}Q_k^a(d)d$ belongs to the class $\chi(p, q, \omega)$, so that it can be realized in terms of ω -periodic state-space realization of the kind (3.1), (3.2). This last step can be accomplished using the minimal realization procedures given in [33]. \triangle

4 Concluding remarks

The invariant polynomial assignment problem has been introduced and solved for linear periodic discrete-time systems. A matrix algebraic approach has been considered and the solution has been provided in terms of a linear periodic discrete-time controller whose associate transfer matrix originates from the solution of a suitable diophantine matrix equation. The synthesis procedure of the controller solving the problem is given in the sufficiency proof of the main theorem.

Acknowledgements

The authors wish to thank proff. G. Conte and A.M. Perdon for their useful discussions and suggestions.

References

- [1] Bittanti S. (1986). Deterministic and stochastic linear periodic systems. *Time Series and Linear Systems*, S.Bittanti Ed., Springer Verlag, 141.
- [2] Grasselli O.M. and S. Longhi (1988). Disturbance localization by measurements feedback for linear periodic discrete-time systems. *Automatica*, **24**, 375-385.
- [3] Davis, J.H. (1972). Stability conditions derived from spectral theory: discrete systems with periodic feedback, *SIAM J. Control*, Vol. 10, No. 1, 1-13.
- [4] Willems, J.L., V. Kučera and P. Brunovsky (1984). On the assignment of invariant factors by time-varying feedback strategies. *Syst. Control Letters*, vol. 5, pp. 75-80.

- [5] Anderson, B.D.O., J.B. Moore (1985). Decentralized control using time-varying feedback, in *Control and Dynamic Systems*, Vol. 22 (C.T. Leondes, Ed.), Academic Press, London, (pp 85-115).
- [6] Kaczorek, T. (1985). Pole placement for linear discrete-time systems by periodic output-feedback, *Syst. Control Lett.*, Vol. 6, 267-269.
- [7] Kaczorek, T. (1987). Invariant factors and pole/variant zero assignments by periodic output-feedback for multivariable systems, *Preprints of the 10th IFAC Congress*, Munich (FRG), Vol. 9, 138-143.
- [8] Khargonekar P.P., K.Poola and A.Tannenbaum (1985). Robust control of linear time-invariant plants using periodic compensators. *IEEE Trans. Aut. Contr.*, AC-30, 1088.
- [9] Grasselli, O.M. , A. Isidori and F. Nicolò (1979). Output regulation of a class of bilinear systems under constant disturbances, *Automatica*, Vol. 15, 189-195.
- [10] Grasselli, O.M., A. Isidori and F. Nicolò (1980). Dead-beat control of discrete-time bilinear systems, *Int. J. of Control*, vol. 32, no. 1, 31-39.
- [11] Graselli, O. M., and S. Longhi (1983). On the stabilization of a class of bilinear systems . *Int. J. Control*, 37, 2, 413-420.
- [12] Kono, M. (1980). Eigenvalue assignment in linear periodic discrete-time systems, *Int. J. Control*, Vol. 32, No. 1, 149-158.
- [13] Ohkawa, F. (1986). Model reference adaptive control system for discrete linear time-varying systems with periodically varying parameters and time delay, *Int. J. Control*, vol. 44, 171-179.
- [14] Zheng, L. (1986). Discrete-time adaptive control for periodically time-varying systems, *Proc. of the IFAC Symposium on Adaptive Systems in Control and Signal Processing*, Lund (Sweden), 131-135.
- [15] Bittanti S., P.Colaneri and G. De Nicolao (1988). The difference periodic Riccati equation for the periodic prediction problem. *IEEE Trans. Aut. Contr.*, AC-33, 706.
- [16] Park, B. and E.I. Verriest (1989). Canonical forms on discrete linear periodically time-varying systems and a control application, *Proc. of the 28th IEEE Confer. on Decision and Control*, Tampa (Florida), 1220-1225.
- [17] Hernandez, V., and A. Urbano (1989). Pole-placement problem for discrete-time linear periodic systems, *Int. J. Control*, vol. 50, 361-371.
- [18] Longhi, S., A. Perdon and G. Conte (1989). Geometric and algebraic structure at infinity of discrete-time linear periodic systems, *Linear Algebra and Applications*, 245-271, pp.245-271.
- [19] Bittanti S., P.Colaneri and G.De Nicolao (1990). An algebraic Riccati equation for the discrete-time periodic prediction problem. *Systems and Control Letters*, 14, 71.
- [20] Colaneri P. (1990). Zero-error regulation for discrete-time linear periodic systems. *Systems and Control Letters*, 15, 377.
- [21] Colaneri P. (1991). Output stabilization via pole-placement of discrete-time linear periodic systems. *IEEE Aut. Contr.*, AC-36, 739.
- [22] Grasselli O.M. and S. Longhi (1991). Robust tracking and regulation of linear periodic discrete-time systems. *Int. J. Contr.*, 54, 613-633.
- [23] Grasselli O.M. and S.Longhi (1991). Pole-placement for nonreachable periodic discrete-time systems. *Math. Contr. Sygn. Systems*, 4, 439-455.
- [24] Dahleh, M.A., P.G. Voulgaris and L.S. Valavani (1992). Optimal and robust controllers for periodic and multirate systems. *IEEE Trans. Autom. Control*, vol. AC-37, no.1, pp 90-99.
- [25] V. Kucěra, *Discrete Linear Control, the Polynomial Equation Approach*, J.Wiley & Sons, Chicester, 1979.
- [26] Kaczorek, T (1993). *Linear Control Systems*, vol. 2, Reseaerch Studies Press LTD, Taunton, England.
- [27] Mayer, R.A., and C.S. Burrus (1975). A unified analysis of multirate and periodically time-varying digital filters. *Trans. Ccts Syst.*, Vol. CSA-22, pp. 162-168.

- [28] Bolzern P., P.Colaneri and R.Scattolini (1986). Zeros of discrete-time linear periodic systems. *IEEE Trans. Aut. Contr.*, **AC-31**, 1057.
- [29] Grasselli O.M. and S. Longhi (1988). Zeros and poles of linear periodic discrete-time systems. *Circuits Systems Signal Process.*, **7**, 361-380.
- [30] Grasselli O.M. and S.Longhi (1991). The geometric approach for linear periodic discrete-time systems, *Linear Algebra and its Applications*, **158**, 27-60.
- [31] Conte, G., A.M. Perdon and S. Longhi (1991). Zero structures at infinity of linear periodic systems, *Circuits Systems Signal Processing*, **10**, pp. 91-100.
- [32] Grasselli O.M. and S.Longhi (1991). Finite zero structure of linear periodic discrete-time systems, *Int. J. Systems Sci.*, **22**, pp. 1785-1806.
- [33] Colaneri, P. and S. Longhi (1993). Discrete-time linear periodic systems: the minimal realization problem, *Proc. of the 12th IFAC World Congress*, Sydney, Australia, vol. 5, pp. 69-74.
- [34] Gohberg, I., M.A. Kaashoek and L. Lerer (1992). Minimality and realization of discrete time-varying systems. *Operator Theory: Advances and Applications*, vol. 56, pp 261-296.